



UNIVERSITI PUTRA MALAYSIA

EFFICACY OF CALCIUM SUPPLEMENTATION AND WEIGHT-BEARING EXERCISE ON BONE MINERAL DENSITY IN POSTMENOPAUSAL CHINESE WOMEN

CHAN YOKE MUN

FPSK (P) 2003 1

**EFFICACY OF CALCIUM SUPPLEMENTATION AND WEIGHT-BEARING
EXERCISE ON BONE MINERAL DENSITY IN POSTMENOPAUSAL
CHINESE WOMEN**

By
CHAN YOKE MUN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirement for the
Degree of Doctor of Philosophy**

May 2003



**This thesis is dedicated to
my parents and my husband, Sing Ziunn**

Abstract of thesis presented to the senate of Universiti Putra Malaysia in fulfillment of the requirements for the degree of Doctor of Philosophy

**EFFICACY OF CALCIUM SUPPLEMENTATION AND WEIGHT-BEARING
EXERCISE ON BONE MINERAL DENSITY IN POSTMENOPAUSAL CHINESE
WOMEN**

By

CHAN YOKE MUN

May 2003

Chairperson: Associate Professor Dr. Zaitun Yassin, Ph.D.

Faculty: Medicine and Health Sciences

The etiology of age-related bone loss is unclear but both habitually low calcium intake and lack of physical activity have been proposed as its determinants. The objective of this two-year randomized controlled trial was to evaluate the efficacy of increasing calcium intake and weight-bearing exercise on bone mineral density (BMD) in postmenopausal Chinese women who were not on hormone replacement therapy.

Bone mineral density at the total body, lumbar spine L₂-L₄, femoral neck, Ward's triangle, trochanter and total hip were measured using dual energy X-ray absorptiometry (DEXA) at six month intervals. Information on sociodemographic background, lifestyle factors and reproductive history were assessed using a validated questionnaire. Calcium intake was evaluated by both the three-day food record and food frequency questionnaire at baseline, 12 and 24 months. Exercise habits were assessed with an exercise diary and Physical Activity Scale for the Elderly (PASE) questionnaire.

The baseline dietary data showed that dietary calcium intake was low among subjects with a mean intake of 444.0 ± 195.4 mg/day. Approximately 30% of the calcium intake was contributed by vegetables, beans and legumes while less than 15% was contributed from milk and dairy products.

The one-way within subjects ANOVA analysis indicated a significant time effect on bone loss for the control group at all skeletal sites. There were significant decreases in BMD at the total body (-0.77%), lumbar spine L₂-L₄ (-0.74%), femoral neck (-1.24%), Ward's triangle (-2.60%), trochanter (-3.17%) and total hip (-2.21%) in the control group at 24 months when compared to baseline. For the calcium group, there was decrease in BMD at Ward's triangle (-0.93%) but the decrease was not significant over time (95% confidence interval [CI]: -2.304 to 0.439; $p > 0.05$). There was virtually no significant bone loss in the calcium-exercise group at all the skeletal sites. There were significant increases in BMD at the femoral neck (+2.62%, $p < 0.05$), trochanter (+4.26%, $p < 0.05$) and total hip (+2.24%, $p < 0.05$). Inter-group comparisons on the changes in BMD over time were computed. Rate of bone loss was significantly higher in the control group as compared to the calcium or calcium-exercise group. There was a significant reduction in rate of bone loss in the calcium-exercise group as compared to calcium group at the femoral neck (+2.62% versus -0.90%), Ward's triangle (+2.17% versus -0.93%), trochanter (+4.26% versus -0.11%) and total hip (+2.24% versus -0.26%). This indicates an additional effect of weight-bearing exercise over calcium supplementation on the hip regions.

In conclusion, increasing calcium intake and weight-bearing exercise were both effective in reducing rate of bone loss in postmenopausal women. The additional effect of weight-bearing exercise on the hip regions may suggest a site-specific effect of weight-bearing exercise on BMD. A major fi

BMD and moderate exercise. This is important as it may be more feasible to encourage the public to engage in moderate exercises such as brisk walking rather than in strenuous exercise. These results support the implementation of a simple public health regimen to retard rate of bone loss by increasing calcium intake and encouraging moderate exercise.

Abstrak tesis yang dikemukakan kepada senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**KEBERKESANAN KALSIUM SUPPLEMENTASI DAN SENAMAN KE ATAS
KETUMPATAN MINERAL TULANG DI KALANGAN WANITA CINA
MENOPAUS**

Oleh

CHAN YOKE MUN

Mei 2003

Pengerusi: Profesor Madya Dr. Zaitun Yassin, Ph.D.

Fakulti: Perubatan dan Sains Kesihatan

Walaupun etiologi kehilangan tulang berikutan penuaan tidak jelas, namun pengambilan kalsium yang rendah dan kekurangan aktiviti fizikal telah dikaitkan sebagai faktor penentunya. Objektif kajian kawalan rawak dua tahun ini adalah untuk menilai keberkesanan peningkatan pengambilan kalsium dan senaman ke atas ketumpatan mineral tulang (KMT) di kalangan wanita Cina posmenopaus yang tidak menerima terapi penggantian hormon.

Ketumpatan mineral tulang jumlah tubuh, lumbar L₂-L₄, pangkal pinggul, Ward, trokanter dan pinggul diukur dengan menggunakan kaedah 'dual energy X-ray absorptionmetry (DEXA)' setiap enam bulan. Maklumat latarbelakang sosiodemografi, faktor gaya hidup dan sejarah reproduktif telah ditentukan dengan penggunaan borang soal selidik yang telah divalidasi. Pengambilan kalsium telah dikaji dengan menggunakan kaedah rekod pengambilan makanan tiga hari dan borang kekerapan

pengambilan makanan yang dijalankan pada permulaan kajian, 12 bulan dan 24 bulan. Senaman telah dinilai dengan menggunakan diari senaman dan borang soal selidik Physical Activity Scale for the Elderly (PASE).

Data pengambilan makanan menunjukkan pengambilan kalsium adalah rendah, dengan purata pengambilan sebanyak 444.0 ± 195.4 mg/hari. Lebih kurang 30% pengambilan kalsium disumbangkan oleh sumber sayur-sayuran dan kekacang sementara susu dan hasil tenusu pula menyumbang kurang daripada 15% daripada jumlah pengambilan kalsium.

Ujian ANOVA satu-hala antara subjek menunjukkan terdapat kesan masa yang signifikan terhadap kehilangan tulang di semua bahagian tulang di kalangan kumpulan subjek kawalan. Sebanyak 0.77% (jumlah tubuh), 0.74% (lumbar), 1.24% (pangkal pinggul), 2.60% (Ward), 3.17% (Trokanter) dan 2.21% (pinggul) kehilangan tulang telah dicatatkan di kalangan kumpulan kawalan pada bulan ke-24 berbanding dengan permulaan kajian. Untuk kumpulan kalsium, walaupun kehilangan tulang sebanyak 0.93% telah dicatatkan pada bahagian Ward tetapi kehilangan itu adalah tidak signifikan (95% jangkal keyakinan: -2.304 kepada 0.439; $p > 0.05$). Untuk kumpulan kalsium-senaman, tidak ada sebarang kehilangan tulang yang dicatatkan di mana-mana bahagian tulang. Terdapat peningkatan KMT pada bahagian pangkal pinggul (+2.62%, $p < 0.05$), trokanter (+4.26%, $p < 0.05$) dan pinggul (+2.24%, $p < 0.05$). Perubahan KMT mengikut masa telah dibandingkan di antara kumpulan kajian. Kadar kehilangan tulang adalah lebih tinggi di kalangan kumpulan kawalan berbanding dengan kumpulan kalsium

ataupun kumpulan kalsium-senaman. Juga, kadar kehilangan tulang yang lebih rendah dicatatkan di kalangan kumpulan kalsium-senaman berbanding kumpulan kalsium di bahagian pinggul (+2.24% berbanding -0.26%), pangkal pinggul (+2.62% berbanding 0.90%), ward (+2.17% berbanding -0.93%) dan trokanter (+4.26% berbanding -0.11%). Ini menunjukkan terdapat kesan tambahan akibat senaman di bahagian pinggul.

Kesimpulannya, peningkatan pengambilan kalsium dan senaman adalah berkesan untuk melambatkan kadar kehilangan tulang di kalangan wanita posmenopaus. Kesan tambahan akibat senaman di kawasan tulang pinggul mencadangkan kesan spesifik setempat senaman ke atas KMT. Perkaitan positif di antara KMT dan senaman sederhana merupakan satu keputusan yang bermakna kerana senaman sederhana seperti berjalan cepat adalah lebih mudah dilakukan berbanding dengan senaman yang lain. Data yang diperolehi menyokong implementasi regimen kesihatan awam untuk mengurangkan kadar kehilangan tulang melalui peningkatan pengambilan kalsium dan senaman sederhana.

ACKNOWLEDGEMENTS

This thesis would not have been possible without the help and expertise of many people. I wish to express my thanks, respect and gratitude to them. My deep appreciation goes to Associate Professor Dr. Zaitun Yassin for her guidance and concern in the preparation of this thesis. Her never-ending support of my work had offered me the impetus to complete my work. I am enormously grateful to my co-supervisors, Associate Professor Dr. Suriah Abdul Rahman, Professor Dr. Chan Siew Pheng and Dr. Mirmalini Kandiah for their careful review of the manuscript, encouragement and many helpful comments.

I would also like to thank the research team, Dr. Winnie Chee Siew Swee, Ms Lee Lai Fun and Mrs Yap Siew Lee for their assistance and encouragement throughout the period of this study. Special thanks to my good friend, Lai Fun, who combed the entire manuscript with her usual care and perception. To my beloved friends, S/N Cheah, Rani, Sophie and Cik Norlisa, thank you for your motivation and moral support.

To those who have helped me in the BMD scanning and biochemical analysis (technicians and laboratory assistants), thank you, for I could not have done it without you all. This thesis has been enriched by the participation of subjects whose continuing support and encouragement are priceless. Finally, I am also grateful for the financial support for this research from the Malaysian Government under the Intensified Research in Priority Research (IRPA) grant no: 06-02-05-9003.

I certify that an Examination Committee met on 5th May 2003 to conduct the final examination of Chan Yoke Mun on her Doctor of Philosophy thesis entitled "Efficacy of Calcium Supplementation and Weight-bearing Exercise on Bone Mineral Density in Postmenopausal Chinese Women" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Khor Geok Lin, Ph.D.

Professor
Department of Nutrition and Health Sciences
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairperson)

Zaitun Yassin, Ph.D.

Associate Professor
Department of Nutrition and Health Sciences
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Suriah Abdul Rahman, Ph.D.

Associate Professor
Program of Food Science
School of Chemical Science and Food Technology
Faculty of Science and Technology
Universiti Kebangsaan Malaysia
(Member)

Mirnalini Kandiah, Ph.D.

Department of Nutrition and Health Sciences
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)



GULAM RUSUL RAHMAT ALI, Ph.D.

Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 28 MAY 2003

The thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:

Zaitun Yassin, Ph.D.

Associate Professor
Department of Nutrition and Health Sciences
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairperson)

Suriah Abdul Rahman, Ph.D.

Associate Professor
Program of Food Science and Nutrition
School of Chemical Science and Food Technology
Faculty of Science and Technology
Universiti Kebangsaan Malaysia
(Member)

Chan Siew Pheng, FRCP

Professor
Department of Medicine
Faculty of Medicine
University Malaya
(Member)

Mirnalini Kandiah, Ph.D.

Department of Nutrition and Health Sciences
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)




AINI IDERIS, Ph.D.

Professor/Dean,
School of Graduate Studies,
Universiti Putra Malaysia.

Date: 14 2 2003

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



CHAN YOKE MUN

Date: 28 MAY 2003

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	ix
APPROVALS	x
DECLARATION	xii
TABLE OF CONTENTS	xiii
LIST OF TABLES	xvi
LIST OF FIGURES	xx
LIST OF ABBREVIATIONS	xxii

CHAPTER

1	INTRODUCTION	1
	1.1 Introduction	1
	1.2 Statement of Problem	2
	1.3 Importance of Study	6
	1.4 Objectives of Study	8
	1.5 Operational Definition	9
2	LITERATURE REVIEW	10
	2.1 Bone and Osteoporosis	10
	2.1.1 Basic Bone Structure and Composition	10
	2.1.2 Bone Architecture	10
	2.1.3 Bone Remodeling Cycle	11
	2.1.4 Markers of Bone Turnover in Relation to Bone Health	13
	2.1.5 Bone and Calcium Homeostatis	20
	2.2 Epidemiology and Outcomes of Osteoporotic Fractures	22
	2.3 Bone Growth, Maintenance and Loss	24
	2.3.1 Skeletal Growth and Maturation	24
	2.3.2 Determinants of Bone Mass	24
	2.4 Risk Factors for Osteopenia and Osteoporosis	31
	2.4.1 Role of Genetic Factors in the Pathogenesis of Osteoporosis	33
	2.4.2 Body Weight, Body Composition and BMD	36
	2.4.3 Nutrition in the Bone Mineral Density Context	39
	2.4.3.1 Calcium Nutriture and Bone Health	40
	2.4.3.2 Vitamin D and Osteoporosis	85
	2.4.3.3 Protein and Bone Health	93
	2.4.3.4 Sodium Intake and Bone Health	100



	2.4.3.5 Vitamin K and Bone Health	101
	2.4.3.6 Micronutrients	103
	2.4.4 The Role of Physical Activity in the Development and Maintenance of BMD	107
	2.4.5 Other Lifestyle Factors	123
2.5	Fall and Osteoporotic Fractures	126
2.6	Bone Mineral Density Measurement	127
	2.6.1 Single Energy Absorptionmetry	128
	2.6.2 Dual-Photon Absorptionmetry	128
	2.6.3 Dual-Energy X-Ray Absorptionmetry (DEXA)	129
3	MATERIALS AND METHODS	136
3.1	General Study Design and Subjects	136
3.2	Screening and Recruitment	138
	3.2.1 Inclusion Criteria	139
	3.2.2 Exclusion Criteria	140
	3.2.3 Recruitment Center	141
3.3	Randomization and Treatment Assignment	143
	3.3.1 Calcium Supplementation Group	144
	3.3.2 Calcium-Exercise Group	145
	3.3.3 Control Group	146
3.4	Data Collection	147
	3.4.1 Socioeconomic and Lifestyle Assessment	147
	3.4.2 Physical Activity Assessment	148
	3.4.3 Dietary Assessment	148
	3.4.4 Calculation of Nutrient Intake	151
	3.4.5 Bone Mineral Density Measurement	153
	3.4.6 Anthropometry Measurements	162
	3.4.7 Laboratory Studies and Biochemical Measurement	165
	3.4.8 Hand Grip Strength Measurement	170
3.5	Statistical Analysis	170
3.6	Limitations of Study	171
3	RESULTS AND DISCUSSION	172
4.1	Randomization and Subject Disposition	172
4.2	Baseline Characteristics of Subjects	174
	4.2.1 Sociodemographic Characteristics	174
	4.2.2 Lifestyle Habits	176
	4.2.3 Reproductive History	178
	4.2.4 Family History of Bone Fractures	180
	4.2.5 Anthropometry Measurements	181
	4.2.6 Bone Mineral Density (BMD)	183
	4.2.7 Dietary Intake Data	189
	4.2.7 Assessment of Physical Activity	206
	4.2.8 Biochemical Data	209
	4.2.9 Determinant of Bone Mineral Density	215

4.2.10	Baseline Comparison Between Different Treatment Group	235
4.3	Effects of Calcium Supplementation and Calcium-Exercise Intervention	238
4.3.1	Changes in Bone Mineral Density	238
4.3.2	Changes in Dietary Intake	253
4.3.3	Changes in Physical Activity	259
4.3.4	Changes in Physical Characteristics	262
4.3.5	Changes in Biochemical Indices	265
4.4	Calcium Tablet and Exercise Acceptance	270
5	CONCLUSION	273
5.1	Summary and Conclusion	273
5.2	Recommendation	278
5.2.1	Population-based Strategies	279
5.2.2	Recommendation for Reviewing Calcium Recommendation	283
5.2.3	Recommendation for Future Research	284
	BIBLIOGRAPHY	286
	APPENDICES	
A	Recommended daily dietary allowance (RDA) Malaysia	336
B	Screening Questionnaire	337
C	Consent Form	340
D	Questionnaire	344
E	Physical Activity Scale for the Elderly (PASE) Questionnaire	351
F	Food Frequency Questionnaire	358
G	Three Day Dietary Record	362
H	DEXA description	367
I	Example of Bone Mineral Densities Analysis	368
J	Calcium Tablet Acceptance	372
K	Calcium Tablet and Exercise Acceptance	373
	VITA	374

LIST OF TABLES

Table		Page
1.1	Percentage distribution of elderly by ethnic group	5
1.2	Operation definitions of variables	9
2.1	Biochemical indices of bone turnover	14
2.2	Risk factors for low bone mass and osteoporosis	32
2.3	Summary of results of randomized trials examining the effect of calcium supplementation on BMD	50
2.4	Recommended daily calcium intake from various sources	66
2.5	Calcium content of some common foods in Malaysia	68
2.6	Calcium content and estimated calcium absorbed of selected food items	75
2.7	Calcium absorption from milk and various salts preparation	82
2.8	Characteristics of RCTs of the efficacy of vitamin D supplementation on rate of bone loss	90
2.9	Recent reports relating bone status to dietary protein intake	99
2.10	Common dietary sources of vitamin K	102
2.11	Currently available BMD measurement techniques	127
2.12	Densitometer precision and minimum duration required to detect BMD change	131
3.1	Measurements/tests conducted for subjects	137
3.2	Summary of numbers of subjects screened and eligible subjects recruited	142
3.3	Estimation of cooking oil and salt used as ingredients	152
3.4	Classification of Body Mass Index according to WHO (1998)	163



4.1	Subject retention at various time-points during the study	173
4.2	Demographic and socioeconomic profile of the subjects	175
4.3	Distribution of respondents according to lifestyle habits	177
4.4	Characteristic of reproductive history	179
4.5	Distribution of respondents according to fracture history	180
4.6	Distribution of respondents according to anthropometric characteristics	181
4.7	Mean \pm s.d. of bone mineral density of subjects	183
4.8	Differences in mean BMD of the lumbar spine among American, Singaporean and Malaysian subjects in this study	186
4.9	Differences in mean BMD of the femoral neck among American, Singaporean and Malaysian subjects in this study	186
4.10	Comparison of BMD with Hong Kong published BMD normogram	188
4.11	Daily energy and nutrient intakes from three-day food record (n=205)	189
4.12	Comparison of mean daily nutrient intakes with Malaysian RDA	191
4.13	Proportion of subjects meeting recommendations for dietary risk	194
4.14	Comparison of nutrient intake with other Malaysian data	195
4.15	Comparison of mean nutrient intake with other studies	197
4.16	Distribution of calcium intake of respondents according to Malaysian RDA and FAO/WHO Recommendation (2002)	198
4.17	Percentage contribution of food sources to dietary calcium in different populations	202
4.18	Distribution of subjects according to patterns of milk consumption	203

4.19	Distribution of subjects according to past physical activity	206
4.20	PASE item weights and contributions to total score	207
4.21	Current pattern of physical activity (n=205)	208
4.22	Mean values of selected biochemical indices of respondents (n=205)	210
4.23	Comparison of vitamin D status in different populations and geographical regions	213
4.24	Correlation of biochemical marker with age and duration of menopause	214
4.25	Mean values of DPD/Creatinine classified by age group and comparison with the Taiji Study	215
4.26	Correlation between BMD and other indices	216
4.27	Proportion of variability of BMD measurements explained by age	217
4.28	Proportion of variability of BMD explained by anthropometric measures	217
4.29	Correlation coefficient of body composition, age and duration of menopause	219
4.30	Correlation of dietary factors, past and current physical activity and milk consumption pattern with BMD	227
4.31	Correlation of biochemical indices with BMD	229
4.32	Multiple regression analysis among selected variables and BMD	232
4.33	Baseline characteristics of subjects (n=172)	236
4.34	Mean percentage change in BMD after 24 months intervention	239
4.35	Mean rate of BMD change in year 1 and year 2 (second-year rate minus first-year rate) in postmenopausal women	248

4.36	Between-Groups difference in BMD changes in postmenopausal women treated with calcium supplementation, calcium-exercise or control for 2 years	249
4.37	Comparison of macronutrient intake over time in the three treatment group and absolute mean changes at 24 months	254
4.38	Comparison of vitamin intake over time and absolute mean changes at 24 months	256
4.39	Comparison of intake of selected minerals over time and absolute mean changes at 24 months	258
4.40	Mean percentage change in PASE score after 24 months intervention	260
4.41	Percentage change in anthropometric indices after 24 months	263
4.42	Selected biochemical indices at baseline, 12 and 24 months	266
4.43	Mean percentage change in 25 (OH)D after 24 months intervention	266
4.44	Consumption practices with calcium supplementation	271

LIST OF FIGURES

Figure		Page
2.1	Determinants of peak bone mass	28
2.2	Mechanism of bone loss in postmenopausal women	30
2.3	Schematic representation of the relationship between various determinants with osteoporosis and osteoporotic-complications	33
3.1	Screening and randomization of subjects into study groups	143
4.1	Randomization and disposition of subjects	173
4.2	Distribution of respondents according to T-score classification by WHO (1994)	185
4.3	Food sources of dietary calcium	201
4.4	Change in total body BMD with time. Results are mean \pm SE. Treatment groups were compared over time using repeated measures ANOVA. Calcium and calcium-exercise were significant different from control group over the duration of study	240
4.5	Change in lumbar spine L ₂ -L ₄ BMD with time. Results are mean \pm SE. Treatment groups were compared over time using repeated measures ANOVA. Calcium and Calcium-exercise were significant different from control group over the duration of study	240
4.6	Change in femoral neck BMD with time. Results are mean \pm SE. Treatment groups were compared over time using repeated measures ANOVA. Calcium-exercise group was significant different from calcium group and control group. Calcium group was significant different from control group over the duration of study	241

4.7	Change in Ward's triangle BMD with time. Results are mean \pm SE. Treatment groups were compared over time using repeated measures ANOVA. Calcium-exercise group was significant different from calcium group and control group. There was no significant different between calcium group and control group over the duration of study	241
4.8	Change in trochanter BMD with time. Results are mean \pm SE. Treatment groups were compared over time using repeated measures ANOVA. Calcium-exercise was significant different from calcium group and control group. Calcium group was significant different from the control group over the duration of study	242
4.9	Change in total hip BMD with time. Results are mean \pm SE. Treatment groups were compared over time using repeated measures ANOVA. Calcium-exercise was significant different from calcium group and control group. Calcium group was significant different from the control group over the duration of study	242
4.10	Relationship between change in femoral neck BMD and duration of brisk walking	261

LIST OF ABBREVIATIONS

BMD	Bone Mineral Density
BMI	Body Mass Index
CI	Confidence Interval
DEXA	Dual Energy X-ray Absorptionmetry
ECF	Extracellular Fluid
NIH	National Institute of Health
PBM	Peak Bone Mass
PTH	Parathyroid Hormone
RDA	Recommended Dietary Allowance
RCT	Randomized Controlled Trial
WHO	World Health Organization
WHR	Waist Hip Ratio

CHAPTER 1

INTRODUCTION

1.1 Introduction

Osteoporosis was defined in 1993 as a systemic disease characterized by low bone mass and microarchitectural deterioration of bone tissue, leading to enhanced bone fragility and increased fracture risk (Consensus Development Conference, 1993). Recently, this definition was modified as a skeletal disorder characterized by compromised bone strength predisposing to an increased risk of fracture (National Institutes of Health, 2000). Bone strength reflects the integration of two main features: bone density and bone quality. In the absence of accurate methods of measuring bone quality, the diagnosis of osteoporosis tends to be made on the basis of low bone density.

The rising incidence of osteoporosis and osteoporotic fractures is becoming a global public health problem. This is based on the recognition that it is a common disease in the developed countries and is likely to become so in the developing countries, where life span is rapidly increasing (Genant et al., 1999).

Primary osteoporosis affects mainly postmenopausal women. Women are at particular risk for fracture, and the data suggests that the incidence in women is twice that in men (Melton, 1988; Hoffenberg et al., 1989; Johnell et al., 1992). This has also been found to be true of Malaysia (Lau et al., 2001a). Above the age of 50 years, one in four women (Melton et al., 1992) and one in eight men (Melton et al., 1992; WHO, 1994) are

believed to have osteoporosis. The National Institutes of Health (NIH) of the United States (1998) pointed out that osteoporosis is about four times more common in women than in men. The higher occurrence of osteoporotic fractures in women is due to several factors. There are attainment of a lower bone mass at the time of maturity (bone mass is approximately 30% higher in men than in women (Consensus Conference, 1984)), women experiencing an accelerated loss of bone after the menopause, the greater likelihood of falls among elderly women (Winner et al., 1989) as well as to women's greater life expectancy (Genant et al., 1999). In total, women lose about 25-30% of the cortical bone and 35-50% of the trabecular bone over their lifetime (Riggs et al., 1981) while men lose about two-thirds of this rate.

1.2 Statement of Problem

Osteoporosis is a chronic and debilitating disease that can influence every facet of a person's life. It has been described as a primary factor contributing to deterioration of quality of life for the elderly (Birge, 1993). It affects more than 75 million people in the United States, Europe and Japan and causes more than 2.3 million fractures annually in the United States and Europe alone (Morii and Genant, 1998).

Postmenopausal bone loss is a major factor in the increasing prevalence of osteoporosis (Aloia et al., 1994). Osteoporosis is usually occult and asymptomatic until a fracture occurs. Thus, fractures are viewed as a complication of osteoporosis, and the relationship between osteoporosis and fractures is analogous to that between high blood pressure and